

Sapozhnikov Ph. V., Kalinina O.Yu. Microphytes of different types of locations from Bunger Hills (Eastern Antarctica). *Voprosy sovremennoi algologii (Issues of modern algology)*. 2019. № 1 (19). С. 57–77. URL: <http://algology.ru/1430>

Таблица 5 (продолжение)

Вид / Местообитание	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	% от 16 местообитаний
<i>Gloeocapsa kuetsingiana</i>	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	12,5
<i>Gloeocapsa sanguinea</i>	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	12,5
<i>Hydrocoryne enteromorphoides</i>	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	12,5
<i>Leptolyngbya laminosa</i>	0	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	12,5
<i>Leptolyngbya nigrescens</i>	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	12,5
<i>Microcoleus vaginatus</i>	0	0	0	0	0	0	0	0	0	1	0	1	0	0	0	0	12,5
<i>Nostoc indistinguendum</i>	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	1	12,5
<i>Nostoc sphaericum</i>	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	0	12,5
<i>Oscillatoria curviceps</i>	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0	12,5
<i>Oscillatoria simplicissima</i>	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	12,5
<i>Phormidium breve</i>	0	0	0	1	0	0	0	1	0	0	0	0	0	0	0	0	12,5
<i>Pseudanabaena catenata</i>	0	0	0	1	0	0	0	0	0	0	0	0	1	0	0	0	12,5
<i>Schizothrix antarctica</i>	0	0	0	0	0	1	0	0	0	0	0	1	0	0	0	0	12,5
<i>Schizothrix simplicior</i>	0	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	12,5
<i>Spirulina corakiana</i>	1	0	0	0	0	0	0	0	0	0	1	0	0	0	0	0	12,5
<i>Tolypothrix distorta</i>	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	12,5
<i>Tolypothrix lanata</i>	0	1	0	0	0	0	0	0	0	0	0	0	1	0	0	0	12,5
<i>Tolypothrix tenuis</i>	0	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	12,5
<i>Tychonema bornetii</i>	0	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	12,5
<i>Anabaena cf. cylindrica</i>	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	6,25
<i>Aphanocapsa cf. muscicola</i>	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	6,25
<i>Aphanocapsa delicatissima</i>	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	6,25
<i>Aphanocapsa elachista</i>	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	6,25
<i>Aphanocapsa grevillei</i>	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Aphanothece clathrata</i>	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	6,25
<i>Asterocapsa cf. divina</i>	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Calothrix cf. braunii</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	6,25
<i>Chroococcus dispersus</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	6,25
<i>Chroococcus subviolaceus</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	6,25

Sapozhnikov Ph. V., Kalinina O.Yu. Microphytes of different types of locations from Bunger Hills (Eastern Antarctica). *Voprosy sovremennoi algologii (Issues of modern algology)*. 2019. № 1 (19). С. 57–77. URL: <http://algology.ru/1430>

Таблица 5 (продолжение)

Вид / Местообитание	1	2	3	4	5	6	8	9	10	11	12	13	14	15	16	17	% от 16 местообитаний
<i>Phormidium acidophilum</i>	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	6,25
<i>Phormidium formosum</i>	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	6,25
<i>Phormidium retzii</i>	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	6,25
<i>Pseudanabaena amphigranulata</i>	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Schizothrix calcicola</i>	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Schizothrix vaginata</i>	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0	6,25
<i>Tolypothrix penicillata</i>	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Xenococcus</i> sp. 1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	6,25
Нaptophyta																	
<i>Ruttnera lamellosa</i>	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	6,25
Ochrophyta																	
<i>Bumilleria</i> cf. <i>exilis</i>	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	6,25
<i>Chrysoreinhardia feldmannii</i>	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0	0	6,25
<i>Pseudopleurochloris</i> sp. 1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0	6,25